INTERNATIONAL STANDARD

ISO 19040-3

First edition
2018-08

Water quality — Determination of the estrogenic potential of water and waste water —

Part 3: In vitro human cell-based reporter gene assay

Qualité de l'eau — Détermination du potentiel oestrogène de l'eau et des eaux résiduaires —

Partie 3: Essai in vitro sur cellules humaines avec gène rapporteur

Reference number
ISO 19040-3:2018(E)

© ISO 2018



COPYRIGHT PROTECTED DOCUMENT

© ISO 2018

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Fax: +41 22 749 09 47
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

 © ISO 2018 – All rights reserved

Contents

Page

Foreword ... **v**

1 Scope ... **1**

2 Normative references ... **1**

3 Terms and definitions ... **1**

4 Interferences ... **4**

5 Principle ... **4**

6 Apparatus and materials ... **4**

7 Reagents, cells and media ... **5**

8 Sampling and samples ... **9**
8.1 General ... 9
8.2 Bottles and material for sampling ... 9
8.3 Bottles and material pre-cleaning ... 9
8.4 Sampling procedure ... 9
8.5 Transport of samples ... 9
8.6 Pretreatment of sample ... 10
8.7 Storage of samples ... 10

9 Procedure ... **10**
9.1 Cell culture maintenance ... 10
9.1.1 Freezing cells ... 10
9.1.2 Starting a new cell culture ... 10
9.1.3 Culturing cells ... 11
9.2 Human cell reporter gene assay test procedure ... 11
9.2.1 Seeding the cells (day 1) ... 11
9.2.2 Preparation of the E2-reference (day 2) ... 11
9.2.3 Preparation of the sample dilutions (day 2) ... 12
9.2.4 Field blank ... 12
9.2.5 Exposing the cells (day 2) ... 12
9.2.6 Harvesting the cells (day 3) ... 13
9.2.7 Measurement of luminescence (day 3) ... 13
9.3 Data analysis ... 13
9.3.1 Calculation of the reporter gene induction ... 13
9.3.2 Calculation of the percentage of maximum response ... 14
9.3.3 Calculation of the dose-response curve ... 14

10 Validity criteria ... **14**
10.1 Validity criteria for the assay ... 14
10.2 Validity criteria for samples ... 15

11 Assessment criteria ... **15**

12 Test report ... **15**

Annex A (informative) **Settings of the luminometer** ... **16**

Annex B (informative) **Plate setup** ... **17**

Annex C (informative) **Bioassay characteristics and details** ... **18**

Annex D (informative) **Test set up for chemicals and extracts** ... **20**

Annex E (informative) **Preparation of dilution series** ... **22**

Annex F (informative) **Performance data** ... **23**

Annex G (informative) **Statistical assessment** ... **33**

Annex H (informative) **Calculation of 17β-estradiol equivalents** ... **34**

© ISO 2018 – All rights reserved

Annex I (informative) **Measurement of the lowest ineffective dilution (LID) of waste water — A simplified evaluation for testing of waste water** **36**

Bibliography **39**

 © ISO 2018 – All rights reserved

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

A list of all parts in the ISO 19040 series can be found on the ISO website.

© ISO 2018 – All rights reserved

Water quality — Determination of the estrogenic potential of water and waste water —

Part 3: In vitro human cell-based reporter gene assay

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices.

IMPORTANT — It is absolutely essential that tests conducted in accordance with this document be carried out by suitably trained staff.

1 Scope

This document specifies a method for the determination of the estrogenic potential of water and waste water by means of a reporter gene assay utilizing stably transfected human cells. This reporter gene assay is based on the activation of the human estrogen receptor alpha.

This method is applicable to:

— fresh water;

— waste water;

— aqueous extracts and leachates;

— eluates of sediments (fresh water);

— pore water;

— aqueous solutions of single substances or of chemical mixtures;

— drinking water;

— the limit of quantification (LOQ) of this method for the direct analysis of water samples is between 0,3 ng/l and 1 ng/l 17β-estradiol equivalents (EEQ) based on the results of the international interlaboratory trial (see Annex F). The upper working range was evaluated [based on the results of the international interlaboratory trial (see Table F.3)] up to a level of 75 ng EEQ/l. Samples showing estrogenic potencies above this threshold have to be diluted for a valid quantification. Extraction and pre concentration of water samples can prove necessary if their estrogenic potential is below the given LOQ.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

© ISO 2018 – All rights reserved

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

— ISO Online browsing platform: available at https://www.iso.org./obp

— IEC Electropedia: available at http://www.electropedia.org/

3.1
culture medium
nutrients presented in a form and phase (liquid or solidified) which support cellular growth

[SOURCE: ISO 6107-6:2004, 24, modified — "cellular" replaces "microbiological"]

3.2
dilution level
D
denominator of the dilution coefficient (using the numerator 1) of a mixture of water or waste water with dilution water as integral number

Note 1 to entry: For undiluted water or waste water, this coefficient per definition is 1→1. The corresponding and smallest possible value of *D* is 1. In this document, the arrow indicates the transition from initial total volume to final total volume.

[SOURCE: ISO 6107-6:2004, 28]

3.3
dilution water
sterile water added to the test sample to prepare a series of defined dilutions

[SOURCE: ISO 20079:2005, 3.7]

3.4
EC_{50}
effective concentration of a compound which causes 50 % of an effect

Note 1 to entry: In the sense of the present document the EC_{50} is the effective concentration of a compound which induces 50 % of the maximal reporter gene activity which can be achieved by this compound.

3.5
extract
test sample after extraction and possible removal of extraction vehicle

3.6
field blank
container prepared in the laboratory, using reagent water or other blank matrix, and sent with the sampling personnel for exposure to the sampling environment to verify possible contamination during sampling

[SOURCE: ISO 11074:2015, 4.5.3]

3.7
induction rate
quotient of the mean value of wells with enhanced reporter gene activity measured on the plates treated with a dose of the test sample or with a positive control, and the mean value of the corresponding wells treated with the negative control using the same cells under identical conditions

[SOURCE: ISO 6107-6:2004, 43, modified — "wells with enhanced reporter gene activity measured" replaces "mutant colonies"; "corresponding wells" replaces "corresponding plates", "quotient" replaces "difference"; "cells" replaces "strain".]

© ISO 2018 – All rights reserved

3.8
limit of quantification
LOQ
lowest value that can be determined with an acceptable level of accuracy and precision

[SOURCE: ISO 15839:2003, 3.18]

3.9
lowest ineffective-dilution value
LID
lowest dilution within a test batch which does not show any effect, i.e. no statistically significant increase in the reporter gene activity compared with the negative control

[SOURCE: ISO 11350:2012, 3.4, modified — "increase in the reporter gene activity" replaces "increase in the number of revertant wells"]

3.10
negative control
dilution water without test sample

[SOURCE: ISO 6107-6:2004, 51]

3.11
passage number
the number of subcultures from cells in a new culture vessel (cell culture flask or micro titer plate)

3.12
reference compound
compound with one or more property values that are sufficiently reproducible and well established to enable the calibration of the measurement method

[SOURCE: ISO 7405:2008, 3.6, modified — "compound" replaces "material"; "the calibration of the measurement method" replaces "use of the material or substance for the calibration of an apparatus, the assessment of a measurement method or for the assignment of values to materials".]

3.13
relative light units
RLU
amount of reporter gene activity as measured by light produces using a luminometer, expressed as relative light units

3.14
reporter gene activity
quantitative activity of a gene attached to the promoter sequence of another gene

3.15
stock culture
frozen culture of cells for the preservation of the characteristics of the cell line

[SOURCE: ISO 21427-2:2006, 13, modified — "the cell line" replaces "V79 cells"]

3.16
subculturing
transfer of part of a cell culture into a new cell culture vessel during cell culture

3.17
test sample
undiluted, diluted or otherwise prepared portion of a sample to be tested, after completion of all preparation steps such as centrifugation, filtration, homogenization, pH adjustment and determination of ionic strength

[SOURCE: ISO 6107-6:2004, 92]

© ISO 2018 – All rights reserved